



## Coverage, Coding, and Payment Guide

**J9023 - Injection, avelumab, 10 mg**

**Please see Important Safety Information on pages 13-15.**

**Click for full [Prescribing Information](#) and [Medication Guide](#), or visit [BAVENCIO.com](#).**

This document is for informational purposes only. It is always the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for products and services rendered. EMD Serono, Inc. does not guarantee coverage and/or reimbursement for BAVENCIO. Coverage, coding, and reimbursement policies vary significantly by payer, patient, and setting of care. Actual coverage and reimbursement decisions are made by individual payers following the receipt of claims. Patients and healthcare professionals should always verify coverage, coding, and reimbursement guidelines on a payer- and patient-specific basis. Please contact CoverOne® at 1-844-8COVER1 (844-826-8371) for support with payer-specific BAVENCIO questions or assistance verifying insurance benefits for a specific patient.

# CoverOne®

CoverOne® provides patient access and reimbursement support services to help eligible patients gain appropriate access to BAVENCIO® (avelumab) Injection 20 mg/mL in the United States.

**We recognize that each patient's situation is different,  
and are dedicated to helping eligible patients one at a time.**



844-8COVER1  
(844-826-8371)  
Monday–Friday  
8:00 AM–5:00 PM ET



800-214-7295



[CoverOne.com](https://CoverOne.com)

## INDICATIONS

BAVENCIO® (avelumab) is indicated for:

- The treatment of adults and pediatric patients 12 years and older with metastatic Merkel cell carcinoma (MCC)
- The maintenance treatment of patients with locally advanced or metastatic urothelial carcinoma (UC) that has not progressed with first-line platinum-containing chemotherapy
- The treatment of patients with locally advanced or metastatic UC who have disease progression during or following platinum-containing chemotherapy, or have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy

BAVENCIO in combination with axitinib is indicated for the first-line treatment of patients with advanced renal cell carcinoma (RCC).

## SELECTED SAFETY INFORMATION

BAVENCIO can cause **severe and fatal immune-mediated adverse reactions** in any organ system or tissue and at any time after starting treatment with a PD-1/PD-L1 blocking antibody, including after discontinuation of treatment.

**Early identification and management of immune-mediated adverse reactions are essential** to ensure safe use of PD-1/PD-L1 blocking antibodies. Monitor patients closely for symptoms and signs that may be clinical manifestations of underlying immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function at baseline and periodically during treatment. In cases of suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

**No dose reduction for BAVENCIO is recommended. For immune-mediated adverse reactions, withhold or permanently discontinue BAVENCIO depending on severity.** In general, withhold BAVENCIO for severe (Grade 3) immune-mediated adverse reactions. Permanently discontinue BAVENCIO for life-threatening (Grade 4) immune-mediated adverse reactions, recurrent severe (Grade 3) immune-mediated reactions that require systemic immunosuppressive treatment, or an inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks of initiating corticosteroids. In general, if BAVENCIO requires interruption or discontinuation, administer systemic corticosteroid therapy (1 to 2 mg/kg/day prednisone or equivalent) until improvement to Grade 1 or less. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reactions are not controlled with corticosteroid therapy. Toxicity management guidelines for adverse reactions that do not necessarily require systemic corticosteroids (eg, endocrinopathies and dermatologic reactions) are discussed in subsequent sections.

Please see Important Safety Information on pages 13-15.  
Click for full [Prescribing Information](#) and [Medication Guide](#), or visit [BAVENCIO.com](https://BAVENCIO.com).

## Table of Contents

<b>Commonly Used Billing Codes for BAVENCIO® (avelumab)</b>	<b>4</b>
National Drug Codes (NDCs)	4
Healthcare Common Procedure Coding System (HCPCS)	4
Current Procedural Terminology (CPT®) Code for Drug Administration Service	5
Revenue Codes—For Use in the Hospital Setting ONLY	5
International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) Diagnosis Codes for Advanced Renal Cell Carcinoma	6
ICD-10-CM Diagnosis Codes for Metastatic Merkel Cell Carcinoma	6
ICD-10-CM Diagnosis Codes for Locally Advanced or Metastatic Urothelial Carcinoma	7
<b>Medicare</b>	<b>8</b>
<b>Medicaid</b>	<b>9</b>
<b>Commercial Payers</b>	<b>9</b>
<b>Sample Claim Forms</b>	<b>10</b>
<b>CoverOne®</b>	<b>12</b>
<b>Important Safety Information</b>	<b>13</b>
<b>References</b>	<b>Back cover</b>

Coverage, coding, and payment for BAVENCIO® (avelumab) Injection 20 mg/mL will vary by payer, plan, setting of care, and patient. The information in this guide provides a general coverage, coding, and payment framework for BAVENCIO across major payer segments. Healthcare professionals should always verify coverage, coding, and payment guidelines on a patient-specific basis.

The CoverOne® program provides patient access and reimbursement support services, and may be reached by calling **1-844-8COVER1 (844-826-8371)**.

## Commonly Used Billing Codes for BAVENCIO

The codes discussed below may be appropriate when billing for BAVENCIO and its administration for the treatment of its FDA-approved indications.

It is always the provider’s responsibility to determine the appropriate healthcare setting and to submit true and correct claims for products and services rendered. EMD Serono, Inc. does not guarantee coverage and/or reimbursement for BAVENCIO. Coverage, coding, and reimbursement policies vary significantly by payer, patient, and setting of care. Actual coverage and reimbursement decisions are made by individual payers following the receipt of claims. Patients and healthcare professionals should always verify coverage, coding, and reimbursement guidelines on a payer- and patient-specific basis.

### National Drug Codes (NDCs)

BAVENCIO NDC (11-digit)	Packaging
<b>44087-3535-01</b>	BAVENCIO is supplied in a single-dose vial of 200 mg/10 mL (20 mg/mL) individually packed

When used in combination with axitinib for first-line advanced RCC, review the full Prescribing Information for axitinib prior to initiation.

### Healthcare Common Procedure Coding System (HCPCS)

HCPCS coding requirements will vary by payer, setting of care, and date of service. Please verify patient-specific insurance benefits to confirm specific coding and billing guidelines for BAVENCIO.

HCPCS <sup>1</sup>	Description	Payer and Setting of Care
<b>J9023</b>	Injection, avelumab, 10 mg	For all payers and settings of care for which HCPCS codes are reported

- 1 unit of J9023 equals 10 mg of avelumab. As a result, 20 units of J9023 equals one 200 mg single-use vial of BAVENCIO® (avelumab) Injection 20 mg/mL, and 80 units equals 800 mg, the recommended dosage for BAVENCIO. Actual units reported will vary by dosage required for each individual patient and any specific billing instructions required by the local payer.
- Medicare claims require the use of the JW modifier (Drug amount discarded/not administered to any patient) when applicable. Effective for dates of service on or after July 1, 2023, Medicare claims require the use of the new JZ modifier (zero drug amount discarded/not administered to any patient) for single use vials when there are no discarded drug amounts.<sup>2</sup> Wastage-reporting policies for payers other than Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.

The recommended dosage of BAVENCIO is 800 mg administered as an intravenous infusion over 60 minutes every 2 weeks until disease progression or unacceptable toxicity. In the event of a Grade 1 or 2 infusion-related reaction, the Dosage Modification section of the Prescribing Information directs providers to interrupt or slow the infusion rate. Please refer to the full BAVENCIO Prescribing Information for complete Dosage and Administration guidelines. When used in combination with axitinib for advanced RCC, review the full Prescribing Information for axitinib prior to initiation.

## Current Procedural Terminology (CPT®)\* Code for Drug Administration Service

CPT®	Description
96413	Chemotherapy administration, intravenous infusion technique; up to 1 hour, single or initial substance/drug
96415	Chemotherapy administration, intravenous infusion technique; each additional hour (list separately in addition to code for primary procedure)

\*CPT Copyright 2023 American Medical Association. All rights reserved. CPT® is a registered trademark of the American Medical Association.

The appropriate CPT® code for the administration of BAVENCIO will depend on the actual service performed. Providers should consult a current CPT® manual and always select the code that accurately describes the administration service based on the specific patient encounter.

## Revenue Codes—For Use in the Hospital Setting ONLY

All hospital claim forms must include a revenue code for each charge line item. The following revenue codes are most relevant for physician-administered drugs.

Code	Code Description
0636	Pharmacy—Drugs Requiring Detailed Coding
025X	Pharmacy

Revenue code 0636 is required on Medicare hospital outpatient claims. For payers other than Medicare, the revenue code may vary. Although some private payers and Medicaid plans accept revenue code 0636, others may require a different revenue code, such as 0250.

Each CPT® code must be reported with a revenue code, which may vary depending on the type of procedure and the cost center in which the procedure is performed.



**Revenue codes are not used in the physician office setting.**

Please see Important Safety Information on pages [13-15](#).  
Click for full [Prescribing Information](#) and [Medication Guide](#), or visit [BAVENCIO.com](#).

## International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) Diagnosis Codes

All hospital and physician office claims must include at least one ICD-10-CM diagnosis code to describe the patient's condition.

### Advanced Renal Cell Carcinoma

ICD-10-CM	Description
C64.1	Malignant neoplasm of right kidney, except renal pelvis
C64.2	Malignant neoplasm of left kidney, except renal pelvis
C64.9	Malignant neoplasm of unspecified kidney, except renal pelvis
C65.1	Malignant neoplasm of right renal pelvis
C65.2	Malignant neoplasm of left renal pelvis
C65.9	Malignant neoplasm of unspecified renal pelvis

### Metastatic Merkel Cell Carcinoma

ICD-10-CM	Description
C4A.0	Merkel cell carcinoma of lip
C4A.10	Merkel cell carcinoma of unspecified eyelid, including canthus
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal
C4A.21	Merkel cell carcinoma of right ear and external auricular canal
C4A.22	Merkel cell carcinoma of left ear and external auricular canal
C4A.30	Merkel cell carcinoma of unspecified part of face
C4A.31	Merkel cell carcinoma of nose
C4A.39	Merkel cell carcinoma of other parts of face
C4A.4	Merkel cell carcinoma of scalp and neck
C4A.51	Merkel cell carcinoma of anal skin
C4A.52	Merkel cell carcinoma of skin of breast
C4A.59	Merkel cell carcinoma of other part of trunk
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder

The ICD-10-CM diagnosis codes listed above are provided only as examples of potentially relevant codes. Providers should consult a current ICD-10-CM manual and always select the most appropriate diagnosis code(s) with the highest level of specificity to describe a patient's actual condition. All diagnosis codes should be supported with adequate documentation in the patient's medical record.

## ICD-10-CM Diagnosis Codes (continued)

### Metastatic Merkel Cell Carcinoma (continued)

ICD-10-CM	Description
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip
C4A.71	Merkel cell carcinoma of right lower limb, including hip
C4A.72	Merkel cell carcinoma of left lower limb, including hip
C4A.8	Merkel cell carcinoma of overlapping sites
C4A.9	Merkel cell carcinoma, unspecified

### Locally Advanced or Metastatic Urothelial Carcinoma

Code	Code Description
C67.0	Malignant neoplasm of trigone of bladder
C67.1	Malignant neoplasm of dome of bladder
C67.2	Malignant neoplasm of lateral wall of bladder
C67.3	Malignant neoplasm of anterior wall of bladder
C67.4	Malignant neoplasm of posterior wall of bladder
C67.5	Malignant neoplasm of bladder neck
C67.6	Malignant neoplasm of ureteric orifice
C67.7	Malignant neoplasm of urachus
C67.8	Malignant neoplasm of overlapping sites of bladder
C67.9	Malignant neoplasm of bladder, unspecified
C65.1	Malignant neoplasm of right renal pelvis
C65.2	Malignant neoplasm of left renal pelvis
C65.9	Malignant neoplasm of unspecified renal pelvis
C66.1	Malignant neoplasm of right ureter
C66.2	Malignant neoplasm of left ureter
C66.9	Malignant neoplasm of unspecified ureter
C68.0	Malignant neoplasm of urethra
C61	Malignant neoplasm of prostate (reporting "urothelial carcinoma of the prostate")
Z85.51	Personal history of malignant neoplasm of bladder
D09.0	Carcinoma in situ of bladder

The ICD-10-CM diagnosis codes listed above are provided only as examples of potentially relevant codes. Providers should consult a current ICD-10-CM manual and always select the most appropriate diagnosis code(s) with the highest level of specificity to describe a patient's actual condition. All diagnosis codes should be supported with adequate documentation in the patient's medical record.

Please see Important Safety Information on pages [13-15](#).  
Click for full [Prescribing Information](#) and [Medication Guide](#), or visit [BAVENCIO.com](#).

## Medicare Fee-For-Service

BAVENCIO® (avelumab) Injection 20 mg/mL is eligible for Medicare Part B fee-for-service coverage when used for an FDA-approved indication and administered “incident to” a physician’s service and when reasonable and necessary for an individual patient.<sup>3</sup>

Medicare Administrative Contractors (MACs) have the discretion to issue Local Coverage Determinations (LCDs) or other written coverage and coding guidance for BAVENCIO. Medicare Part B claims for BAVENCIO in the physician office and hospital outpatient settings should be billed with HCPCS code J9023 (Injection, avelumab, 10 mg).

Medicare claims require the use of the JW modifier (drug amount discarded/not administered to any patient) when applicable. Effective for dates of service on or after July 1, 2023, Medicare claims require the use of the new JZ modifier (zero drug amount discarded/not administered to any patient) for single use vials when there are no discarded drug amounts.<sup>2</sup>

For Medicare hospital outpatient prospective payment system (OPPS) claims, certain 340B providers must use one of the following modifiers with HCPCS code J9023 to indicate that BAVENCIO was acquired through the 340B Drug Pricing Program (when applicable):

- “JG” modifier (Drug or biological acquired with 340B drug pricing program discount, reported for informational purposes) - NOTE: Should be used by 340B hospitals paid under OPPS except for rural sole community hospital, children’s hospitals, and PPS-exempt cancer hospitals
- “TB” modifier (Drug or biological acquired with 340B drug pricing program discount, reported for informational purposes for select entities) NOTE: Should only be used by rural sole community hospitals, children’s hospitals, and PPS-exempt cancer hospitals

Note: In 2023, use of the JG and TB modifiers is required only for informational purposes and will not result in a payment reduction.<sup>4</sup> 340B modifier requirements are subject to change at the direction of CMS. These requirements do not apply to physician offices or non-Medicare payers.

Medicare Part B payment for BAVENCIO in the physician office setting and OPPS hospital outpatient setting is currently based on Average Sales Price + 6%.<sup>5,6</sup>

Medicare payments are subject to 2% sequestration reductions as required under the Budget Control Act of 2011 (P.L. 112-25). CMS updates Part B drug payment rates on a quarterly basis.<sup>7</sup>

Please contact your local MAC or the CoverOne® program to verify specific BAVENCIO coverage, coding, and payment details for your geographic area and clinical care setting. The CoverOne program is available to assist patients and providers and may be reached by calling 1-844-8COVER1 (844-826-8371).



## Medicare Advantage

Medicare Advantage plans must provide access to all Part A and Part B benefits that would be available in Original Medicare Fee-for-Service.<sup>8</sup> However, Medicare Advantage plans may require prior authorization for BAVENCIO. Additionally, starting in 2019, Medicare Advantage plans have the discretion to use step therapy limits for Part B drugs under certain circumstances. This CMS guidance recognizes that MA plans may apply step therapy for Part B drugs as long as it does not create an undue access barrier for beneficiaries.<sup>a</sup> Contact CoverOne or the Medicare Advantage plan directly to confirm plan-specific coverage requirements.

<sup>a</sup>CMS Memo to MA Plans, Prior Authorization and Step Therapy for Part B Drugs in Medicare Advantage, August 7, 2018.

### IMPORTANT NOTE:

Many Medicare enrollees receive coverage through a Medicare Advantage plan. Reimbursement policies under these plans may differ from those of traditional Medicare and may vary from plan to plan.

## Medicaid

Medicaid coverage, coding, and payment policies will vary by state Medicaid program and should be verified for each particular state. In addition, some Medicaid plans may require prior authorization for BAVENCIO® (avelumab) Injection 20 mg/mL.

HCPCS code J9023 (Injection, avelumab, 10 mg) should be used to bill for BAVENCIO in the physician office and hospital outpatient settings, unless otherwise directed by the state Medicaid program. Providers are also required to report the 11-digit NDC on Medicaid claims for all provider-administered drugs.

Medicaid programs use a variety of payment methodologies for drugs and biologicals in the physician office setting, including Average Wholesale Price (AWP),<sup>b</sup> ASP, state-specific fee schedules, or percentage of charges.

Medicaid programs use a variety of payment methodologies for drugs and biologicals in the hospital outpatient setting, including AWP, ASP, ambulatory payment groups, state-specific fee schedules, preset per-diem/per-visit rates, or percentage of charges.

There is significant variation in Medicaid payment amounts among states. Providers should verify payment rates for their state Medicaid program.

The CoverOne® program is available to assist patients and providers in verifying state Medicaid policies for BAVENCIO and may be reached by calling **1-844-8COVER1 (844-826-8371)**.

### IMPORTANT NOTE:

Many Medicaid enrollees receive coverage through Medicaid managed care plans. Coverage and payment policies under these plans may differ from those of traditional Medicaid and may vary from plan to plan.

<sup>b</sup>AWP is set by certain third-party pricing services. EMD Serono, Inc. does not set AWP.

## Commercial Payers

Coverage, coding, and payment guidelines vary significantly by private/commercial payer, plan, setting of care, and patient. Additionally, benefits may also depend on the specific contract terms that a provider negotiates with a given plan. Patients and providers should verify patient-specific benefits to determine if BAVENCIO is covered under each specific plan and if prior authorization is required.

HCPCS code J9023 (Injection, avelumab, 10 mg) should be used to bill for BAVENCIO in the physician and hospital outpatient settings, unless otherwise directed by the private/commercial payer. Many private payers also require the reporting of 11-digit NDC codes in addition to HCPCS codes.

Private insurers use a variety of payment methodologies for drugs and biologicals administered in the physician office and hospital outpatient settings, including AWP, ASP, invoice, or percentage of charges.

The exact payment mechanism used by a specific payer usually depends on the provider's contractual agreement with that payer.

The CoverOne program is available to assist patients and providers in verifying private insurance benefits and may be reached by calling **1-844-8COVER1 (844-826-8371)**.

Please see Important Safety Information on pages **13-15**.  
Click for full [Prescribing Information](#) and [Medication Guide](#), or visit [BAVENCIO.com](https://www.bavencio.com).

## HEALTH INSURANCE CLAIM FORM

## PHYSICIAN OFFICE

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

PICA		PICA	
1. MEDICARE (Medicare#)	MEDICAID (Medicaid#)	TRICARE (ID#/DoD#)	CHAMPVA (Member ID#)
2. PATIENT'S NAME (Last Name, First Name, Middle Initial)		3. PATIENT'S BIRTH DATE MM DD YY	
5. PATIENT'S ADDRESS (No., Street)		6. PATIENT RELATIONSHIP TO INSURED Self <input type="checkbox"/> Spouse <input type="checkbox"/> Child <input type="checkbox"/>	
CITY	STATE	8. RESERVED FOR NUCC USE	
ZIP CODE	TELEPHONE (Include Area Code)		
9. OTHER INSURED'S NAME (Last Name, First Name, Middle Initial)		10. IS PATIENT'S CONDITION RELATED TO PREVIOUS CLAIM?	
a. OTHER INSURED'S POLICY OR GROUP NUMBER		a. EMPLOYMENT? (Current or Previous)	
b. RESERVED FOR NUCC USE		b. AUTO ACCIDENT?	
c. RESERVED FOR NUCC USE		c. OTHER ACCIDENT?	
d. INSURANCE PLAN NAME OR PROGRAM NAME		10d. CLAIM CODES (Designated by NUC)	

**BOX 19 (Electronic claim form: Loop 2400)**  
 Select payers may require specific information in this remarks field, such as NDC code or other information as specified by the payer.

**BOX 21 (Electronic claim form: Loop 2300, Segment HI)**  
 Enter the appropriate primary diagnosis code from the patient's medical record in Item 21A. For example, renal cell carcinoma is most commonly described by ICD-10-CM series C64, and C65. Urothelial carcinoma is most commonly described by ICD-10-CM series C67, C66, C65, and C68.0. Merkel cell carcinoma is described by ICD-10-CM series C4A. Use Item 21B-L to report any applicable secondary diagnoses.  
 Example ICD-10-CM code shown; report the ICD-10-CM code(s) that reflect the patient's actual condition. The reported ICD-10-CM code(s) should reflect the highest level of specificity.

**BOX 24A-B (Electronic claim form: BOX 24A: Loop 2400, Segment DTP BOX 24B: Loop 2300/2400, Segment CLM/SV)**  
 Enter the date of service and the appropriate place of service code.  
 Note: If NDC reporting is required (for example, for Medicaid and many commercial payers), enter the NDC information for BAVENCIO in the shaded portion of BOX 24A above the date of service. Check with the payer to determine the proper format for NDC reporting.

**BOX 24E (Electronic claim form: Loop 2400, Segment SV)**  
 Specify the diagnosis from Box 21 that corresponds to the product or procedure listed in Box 24D.

**BOX 24F (Electronic claim form: Loop 2400, Segment SV102)**  
 Enter the total charge for each line item.

**BOX 24D (Electronic claim form: Loop 2400, Segment SV)**  
 Enter the appropriate HCPCS code for BAVENCIO:  
**J9023 - Injection, avelumab, 10 mg<sup>a</sup>**  
 If applicable, discarded product should be reported on a separate line with the HCPCS code and JW modifier. Effective for dates of service on or after July 1, 2023, Medicare claims require the use of the new JZ modifier for single use vials when there are no discarded drug amounts.<sup>b</sup>  
 Enter the appropriate CPT code for the administration service. For example, a 60-minute chemotherapy IV infusion would be reported with CPT code 96413 - *Chemotherapy administration, intravenous infusion technique; up to 1 hour, single or initial substance/drug.*

**BOX 24G (Electronic claim form: Loop 2400, Segment SV)**  
 Enter the number of service units for each line item. For J9023, each unit corresponds to 10 mg of BAVENCIO. One 200 mg single-dose vial would be reported with 20 units. Four 200 mg single-dose vials (800 mg = recommended dosage) would be reported with 80 units.  
 If applicable for the patient encounter, and required by the payer, enter the number of units discarded, corresponding with the line item with the JW modifier.

LINE	DATE(S) OF SERVICE	PROCEDURES, SERVICES, OR SUPPLIES	CHARGE	UNITS	QUAL.	PROVIDER ID. #
1	MM DD YY	J9023				
2	MM DD YY	96413				

Source: Centers for Medicare and Medicaid Services, 2023 Alpha-Numeric Healthcare Common Procedure Coding System File, October 2023  
 \*CMS, CY 2023 Medicare Physician Fee Schedule Final Rule (public display version), November 2, 2022, Page 939.

<sup>a</sup>It is always the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for products and services rendered. EMD Serono, Inc. does not guarantee coverage and/or reimbursement for BAVENCIO. Coverage, coding, and reimbursement policies vary significantly by payer, patient, and setting of care.

Actual coverage and reimbursement decisions are made by individual payers following receipt of claims. Patients and healthcare professionals should always verify coverage, coding, and reimbursement guidelines on a payer- and patient-specific basis.

Please contact CoverOne® at 1-844-8COVER1 (844-826-8371) for support with payer-specific BAVENCIO questions, or assistance verifying insurance benefits for a specific patient.

## HOSPITAL OUTPATIENT DEPARTMENT

## BOX 42

(837I electronic claim: Loop 2400, Segment type SV2)

List the appropriate revenue code for BAVENCIO.

Revenue code 0636 is required by Medicare. For payers other than Medicare, the revenue code may vary; although some private payers and Medicaid plans accept revenue code 0636, others may require a different revenue code, such as 0250.

Enter an appropriate revenue code for the administration service based on the cost center in which the service is performed.

0636

J9023  
96413

## BOX 43

Enter the corresponding description for the revenue code listed in Box 42.

Note: If NDC reporting is required (for example, for Medicaid and many commercial payers), enter the NDC information for BAVENCIO in the shaded portion of BOX 43. Check with the payer to determine the proper format for NDC reporting.

## BOX 44 (837I electronic claim: Loop 2400, Segment type SV2, Service line SV201)

Enter the appropriate HCPCS code for BAVENCIO:

**J9023 - Injection, avelumab, 10 mg<sup>a</sup>**

If applicable, discarded product should be reported on a separate line with the HCPCS code and JW modifier. Effective for dates of service on or after July 1, 2023, Medicare claims require the use of the new JZ modifier for single use vials when there are no discarded drug amounts.<sup>b</sup>

For Medicare OPPS claims, certain 340B providers must use the applicable modifier (JG or TB, depending on the type of facility) to indicate that BAVENCIO was acquired through the 340B Drug Pricing Program (when applicable); see page 8 for additional information. Medicare 340B modifier requirements are subject to change at the direction of CMS.

Enter the appropriate CPT code for the administration service. For example, a 60-minute chemotherapy IV infusion would be reported with CPT code 96413 - *Chemotherapy administration, intravenous infusion technique; up to 1 hour, single or initial substance/drug.*

## BOX 45

(837I electronic claim: Loop 2400, Segment type DTP)

Enter date of service.

## BOX 46 (837I electronic claim: Loop 2400, Segment type SV2, Service line SV205)

Enter the number of service units for each line item. For J9023, each unit corresponds to 10 mg of BAVENCIO. One 200 mg single-dose vial would be reported with 20 units. Four 200 mg single-dose vials (800 mg = recommended dosage) would be reported with 80 units.

If applicable for the patient encounter, and required by the payer, enter the number of units discarded, corresponding with the line item with the JW modifier.

## BOX 47 (837I electronic claim: Loop 2300, Segment type CLM, Claim Information CLM02)

Enter the total charge for each line item.

## BOX 67 (837I electronic claim: Loop 2300, Segment type HI, Diagnosis Information H101-2)

Enter the primary diagnosis code. For example, renal cell carcinoma is most commonly described by ICD-10-CM series C64, and C65. Urothelial carcinoma is most commonly described by ICD-10-CM series C67, C66, C65, and C68.0. Merkel cell carcinoma is described by ICD-10-CM series C4A. Use fields A-Q to report any applicable secondary diagnoses.

Example ICD-10-CM code shown; report the ICD-10-CM code(s) that reflect the patient's actual condition. The reported ICD-10-CM code(s) should reflect the highest level of specificity.

## BOX 80 (837I electronic claim: Loop 2300, Segment type NTE)

Select payers may require specific information in this remarks field.

C67.X

68

69

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

<sup>a</sup>Centers for Medicare and Medicaid Services (CMS), 2023 Alpha-Numeric Healthcare Common Procedure Coding System File, October 2023.<sup>b</sup>CMS, CY 2023 Medicare Physician Fee Schedule Final Rule (public display version), November 2, 2022, Page 939.

It is always the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for products and services rendered. EMD Serono, Inc. does not guarantee coverage and/or reimbursement for BAVENCIO. Coverage, coding, and reimbursement policies vary significantly by payer, patient, and setting of care. Actual coverage and reimbursement decisions are made by individual payers following the receipt of claims. Patients and healthcare professionals should always verify coverage, coding, and reimbursement guidelines on a payer- and patient-specific basis.

Please contact CoverOne® at 844-8COVER1 (844-826-8371) for support with payer-specific BAVENCIO questions or assistance verifying insurance benefits for a specific patient.



**CoverOne®** is a patient access and reimbursement support program available to help eligible patients gain appropriate access to BAVENCIO® (avelumab) Injection 20 mg/mL in the United States. CoverOne offers the following:

## Reimbursement Support

- Insurance Benefit Verification
- Prior Authorization Assistance
- Information on Relevant Billing Codes for BAVENCIO
  - HCPCS, CPT®, ICD-10-CM, NDC
- Denied/Underpaid Claims Assistance
- Payer Research (non-patient specific): Medicare, Private Payers, State Medicaid

## Patient Assistance Program

CoverOne includes a patient assistance program that provides BAVENCIO at no charge for eligible patients who meet certain income, insurance (ie, uninsured), and residency eligibility criteria.

- To determine patient eligibility, providers should complete and fax a CoverOne Enrollment Form **prior to treatment** or submit an online enrollment form at [CoverOne.com](https://CoverOne.com)
- Patient assistance is not applied retroactively

## Co-pay Assistance Program

CoverOne provides co-pay assistance for **privately insured** BAVENCIO patients with co-pay/co-insurance responsibilities who meet the program eligibility criteria.

- Privately insured patients may apply for assistance through the CoverOne Co-pay Program by faxing a completed CoverOne Enrollment Form, or by submitting an online enrollment form at [CoverOne.com](https://CoverOne.com)
  - **Government-insured patients, including beneficiaries of Medicare and Medicaid or other federal or state healthcare programs, are not eligible for the CoverOne Co-pay Assistance Program**
- Limits, terms, and conditions apply. Full terms and conditions for co-pay assistance can be found at [CoverOne.com](https://CoverOne.com)
- CoverOne will notify you of the eligibility determination as soon as possible
- Enrolled patients may be eligible to pay as little as a \$0 co-pay for each treatment for BAVENCIO, up to a maximum co-pay assistance amount of \$30,000 per year
  - Once the annual co-pay assistance limit is reached, enrolled patients are responsible for paying all co-pays and any balance not covered by CoverOne
- For enrolled patients, disbursement of co-pay assistance funds occurs after the patient has received treatment in an outpatient setting, and an Explanation of Benefits (EOB) showing a separately payable BAVENCIO claim eligible for co-pay assistance is sent to CoverOne
- EMD Serono reserves the right to rescind, revoke, or amend the program without notice at any time

PHONE: **844-8COVER1** (844-826-8371) | FAX: **800-214-7295**

[CoverOne.com](https://CoverOne.com) | Monday-Friday, 8:00 AM-5:00 PM ET

When BAVENCIO is used in combination with axitinib for first-line advanced RCC, review the full Prescribing Information for axitinib prior to initiation.

## INDICATIONS and IMPORTANT SAFETY INFORMATION for BAVENCIO® (avelumab)

### INDICATIONS

BAVENCIO® (avelumab) is indicated for:

- The treatment of adults and pediatric patients 12 years and older with metastatic Merkel cell carcinoma (MCC)
- The maintenance treatment of patients with locally advanced or metastatic urothelial carcinoma (UC) that has not progressed with first-line platinum-containing chemotherapy
- The treatment of patients with locally advanced or metastatic UC who have disease progression during or following platinum-containing chemotherapy or have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy

BAVENCIO® (avelumab) in combination with axitinib is indicated for the first-line treatment of patients with advanced renal cell carcinoma (RCC).

### IMPORTANT SAFETY INFORMATION

BAVENCIO can cause **severe and fatal immune-mediated adverse reactions** in any organ system or tissue and at any time after starting treatment with a PD-1/PD-L1 blocking antibody, including after discontinuation of treatment.

**Early identification and management of immune-mediated adverse reactions are essential** to ensure safe use of PD-1/PD-L1 blocking antibodies. Monitor patients closely for symptoms and signs that may be clinical manifestations of underlying immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function at baseline and periodically during treatment. In cases of suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

**No dose reduction for BAVENCIO is recommended. For immune-mediated adverse reactions, withhold or permanently discontinue BAVENCIO depending on severity.** In general, withhold BAVENCIO for severe (Grade 3) immune-mediated adverse reactions. Permanently discontinue BAVENCIO for life-threatening (Grade 4) immune-mediated adverse reactions, recurrent severe (Grade 3) immune-mediated reactions that require systemic immunosuppressive treatment, or an inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks of initiating corticosteroids. In general, if BAVENCIO requires interruption or discontinuation, administer systemic corticosteroid therapy (1 to 2 mg/kg/day prednisone or equivalent) until improvement to Grade 1 or less. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reactions are not controlled with corticosteroid therapy. Toxicity management guidelines for adverse reactions that do not necessarily require systemic corticosteroids (eg, endocrinopathies and dermatologic reactions) are discussed in subsequent sections.

BAVENCIO can cause **immune-mediated pneumonitis**. Withhold BAVENCIO for Grade 2, and permanently discontinue for Grade 3 or Grade 4 pneumonitis. Immune-mediated pneumonitis occurred in 1.1% (21/1854) of patients, including fatal (0.1%), Grade 4 (0.1%), Grade 3 (0.3%), and Grade 2 (0.6%) adverse reactions. Systemic corticosteroids were required in all (21/21) patients with pneumonitis.

BAVENCIO can cause **immune-mediated colitis**. The primary component of immune-mediated colitis consisted of diarrhea. Cytomegalovirus infection/reactivation has been reported in patients with corticosteroid-refractory immune-mediated colitis. In cases of corticosteroid-refractory colitis, consider repeating infectious workup to exclude alternative etiologies. Withhold BAVENCIO for Grade 2 or Grade 3, and permanently discontinue for Grade 4 colitis. Immune-mediated colitis occurred in 1.5% (27/1854) of patients, including Grade 3 (0.4%) and Grade 2 (0.8%) adverse reactions. Systemic corticosteroids were required in all (27/27) patients with colitis.

BAVENCIO can cause **hepatotoxicity and immune-mediated hepatitis**. Withhold or permanently discontinue BAVENCIO based on tumor involvement of the liver and severity of aspartate aminotransferase (AST), alanine aminotransferase (ALT), or total bilirubin elevation. Immune-mediated hepatitis occurred with BAVENCIO as a single agent in 1.1% (20/1854) of patients, including fatal (0.1%), Grade 3 (0.8%), and Grade 2 (0.2%) adverse reactions. Systemic corticosteroids were required in all (20/20) patients with hepatitis.

BAVENCIO **in combination with axitinib** can cause **hepatotoxicity** with higher than expected frequencies of Grade 3 and 4 ALT and AST elevation compared to BAVENCIO alone. Consider more frequent monitoring of liver enzymes as compared to when the drugs are used as monotherapy. Withhold or permanently discontinue both BAVENCIO and axitinib based on severity of AST, ALT, or total bilirubin elevation, and consider administering corticosteroids as needed. Consider rechallenge with BAVENCIO or axitinib, or sequential rechallenge with both BAVENCIO and axitinib, after recovery. In patients treated with BAVENCIO in combination with axitinib in the advanced RCC trials, increased ALT and increased AST were reported in 9% (Grade 3) and 7% (Grade 4) of patients. Immune-mediated hepatitis was reported in 7% of patients including 4.9% with Grade 3 or 4 immune-mediated hepatitis. Thirty-four patients were treated with corticosteroids and one patient was treated with a non-steroidal immunosuppressant.



## IMPORTANT SAFETY INFORMATION for BAVENCIO® (avelumab) (cont'd)

BAVENCIO can cause primary or secondary **immune-mediated adrenal insufficiency**. For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment, including hormone replacement, as clinically indicated. Withhold BAVENCIO for Grade 3 or Grade 4 endocrinopathies until clinically stable or permanently discontinue depending on severity. Immune-mediated adrenal insufficiency occurred in 0.6% (11/1854) of patients, including Grade 3 (0.1%) and Grade 2 (0.4%) adverse reactions. Systemic corticosteroids were required in all (11/11) patients with adrenal insufficiency.

BAVENCIO can cause **immune-mediated hypophysitis**. Hypophysitis can present with acute symptoms associated with mass effect such as headache, photophobia, or visual field defects. Hypophysitis can cause hypopituitarism. Initiate hormone replacement, as clinically indicated. Withhold BAVENCIO for Grade 3 or Grade 4 endocrinopathies until clinically stable or permanently discontinue depending on severity. Immune-mediated pituitary disorders occurred in 0.1% (1/1854) of patients, which was a Grade 2 (0.1%) adverse reaction.

BAVENCIO can cause **immune-mediated thyroid disorders**. Thyroiditis can present with or without endocrinopathy. Hypothyroidism can follow hyperthyroidism. Initiate hormone replacement for hypothyroidism or institute medical management of hyperthyroidism, as clinically indicated. Withhold BAVENCIO for Grade 3 or Grade 4 endocrinopathies until clinically stable or permanently discontinue depending on severity. Thyroiditis occurred in 0.2% (4/1854) of patients, including Grade 2 (0.1%) adverse reactions. Hyperthyroidism occurred in 0.4% (8/1854) of patients, including Grade 2 (0.3%) adverse reactions. Systemic corticosteroids were required in 25% (2/8) of patients with hyperthyroidism. Hypothyroidism occurred in 5% (97/1854) of patients, including Grade 3 (0.2%) and Grade 2 (3.6%) adverse reactions. Systemic corticosteroids were required in 6% (6/97) of patients with hypothyroidism.

BAVENCIO can cause **immune-mediated type I diabetes mellitus**, which can present with diabetic ketoacidosis. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold BAVENCIO for Grade 3 or Grade 4 endocrinopathies until clinically stable or permanently discontinue depending on severity. Immune-mediated type I diabetes mellitus occurred in 0.2% (3/1854) of patients, including Grade 3 (0.2%) adverse reactions.

BAVENCIO can cause **immune-mediated nephritis with renal dysfunction**. Withhold BAVENCIO for Grade 2 or Grade 3, and permanently discontinue for Grade 4 increased blood creatinine. Immune-mediated nephritis with renal dysfunction occurred in 0.1% (2/1854) of patients, including Grade 3 (0.1%) and Grade 2 (0.1%) adverse reactions. Systemic corticosteroids were required in all (2/2) patients with nephritis with renal dysfunction.

BAVENCIO can cause **immune-mediated dermatologic adverse reactions**, including rash or dermatitis. Exfoliative dermatitis including Stevens Johnson Syndrome (SJS), drug rash with eosinophilia and systemic symptoms (DRESS), and toxic epidermal necrolysis (TEN), has occurred with PD-1/PD-L1 blocking antibodies. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-exfoliative rashes. Withhold BAVENCIO for suspected and permanently discontinue for confirmed SJS, TEN, or DRESS. Immune-mediated dermatologic adverse reactions occurred in 6% (108/1854) of patients, including Grade 3 (0.1%) and Grade 2 (1.9%) adverse reactions. Systemic corticosteroids were required in 25% (27/108) of patients with dermatologic adverse reactions.

BAVENCIO can result in **other immune-mediated adverse reactions**. Other clinically significant immune-mediated adverse reactions occurred at an incidence of <1% in patients who received BAVENCIO or were reported with the use of other PD-1/PD-L1 blocking antibodies. For **myocarditis**, permanently discontinue BAVENCIO for Grade 2, Grade 3, or Grade 4. For **neurological toxicities**, withhold BAVENCIO for Grade 2 and permanently discontinue for Grade 3 or Grade 4.

BAVENCIO can cause severe or life-threatening **infusion-related reactions**. Premedicate patients with an antihistamine and acetaminophen prior to the first 4 infusions and for subsequent infusions based upon clinical judgment and presence/severity of prior infusion reactions. Monitor patients for signs and symptoms of infusion-related reactions, including pyrexia, chills, flushing, hypotension, dyspnea, wheezing, back pain, abdominal pain, and urticaria. Interrupt or slow the rate of infusion for Grade 1 or Grade 2 infusion-related reactions. Permanently discontinue BAVENCIO for Grade 3 or Grade 4 infusion-related reactions. Infusion-related reactions occurred in 26% of patients, including three (0.2%) Grade 4 and ten (0.5%) Grade 3 infusion-related reactions. Eleven (85%) of the 13 patients with Grade ≥3 reactions were treated with intravenous corticosteroids.

Fatal and other serious **complications of allogeneic hematopoietic stem cell transplantation (HSCT)** can occur in patients who receive HSCT before or after being treated with a PD-1/PD-L1 blocking antibody. Follow patients closely for evidence of transplant-related complications and intervene promptly. Consider the benefit versus risks of treatment with a PD-1/PD-L1 blocking antibody prior to or after an allogeneic HSCT.

BAVENCIO in combination with axitinib can cause **major adverse cardiovascular events (MACE)** including severe and fatal events. Consider

## IMPORTANT SAFETY INFORMATION for BAVENCIO® (avelumab) (cont'd)

baseline and periodic evaluations of left ventricular ejection fraction. Monitor for signs and symptoms of cardiovascular events. Optimize management of cardiovascular risk factors, such as hypertension, diabetes, or dyslipidemia. Permanently discontinue BAVENCIO and axitinib for Grade 3-4 cardiovascular events. MACE occurred in 7% of patients with advanced RCC treated with BAVENCIO in combination with axitinib compared to 3.4% treated with sunitinib in a randomized trial. These events included death due to cardiac events (1.4%), Grade 3-4 myocardial infarction (2.8%), and Grade 3-4 congestive heart failure (1.8%).

BAVENCIO can cause **fetal harm** when administered to a pregnant woman. Advise patients of the potential risk to a fetus including the risk of fetal death. Advise females of childbearing potential to use effective contraception during treatment with BAVENCIO and for at least 1 month after the last dose of BAVENCIO. It is not known whether BAVENCIO is excreted in human milk. Advise a lactating woman **not to breastfeed** during treatment and for at least 1 month after the last dose of BAVENCIO due to the potential for serious adverse reactions in breastfed infants.

**The most common adverse reactions** (all grades,  $\geq 20\%$ ) in patients with **metastatic Merkel cell carcinoma (MCC)** were fatigue (47%), musculoskeletal pain (29%), infusion-related reaction (26%), rash (25%), nausea (23%), constipation (22%), cough (22%), and diarrhea (21%).

**Laboratory abnormalities** worsening from baseline (all grades,  $\geq 20\%$ ) in patients with **metastatic MCC** were decreased lymphocyte count (51%), decreased hemoglobin (40%), increased aspartate aminotransferase (31%), decreased platelet count (23%), increased alanine aminotransferase (22%), and increased lipase (21%).

A **fatal adverse reaction** (sepsis) occurred in one (0.3%) patient with **locally advanced or metastatic urothelial carcinoma (UC)** receiving BAVENCIO + best supportive care (BSC) as first-line maintenance treatment. In patients with previously treated locally advanced or metastatic UC, fourteen patients (6%) who were treated with BAVENCIO experienced either pneumonitis, respiratory failure, sepsis/urosepsis, cerebrovascular accident, or gastrointestinal adverse events, which led to death.

**The most common adverse reactions** (all grades,  $\geq 20\%$ ) in patients with **locally advanced or metastatic UC** receiving BAVENCIO + BSC (vs BSC alone) as first-line maintenance treatment were fatigue (35% vs 13%), musculoskeletal pain (24% vs 15%), urinary tract infection (20% vs 11%), and rash (20% vs 2.3%). In patients with previously treated locally advanced or metastatic UC receiving BAVENCIO, the most common adverse reactions (all grades,  $\geq 20\%$ ) were fatigue, infusion-related reaction, musculoskeletal pain, nausea, decreased appetite, and urinary tract infection.

**Selected laboratory abnormalities** worsening from baseline (all grades,  $\geq 20\%$ ) in patients with **locally advanced or metastatic UC** receiving BAVENCIO + BSC (vs BSC alone) as first-line maintenance treatment were blood triglycerides increased (34% vs 28%), alkaline phosphatase increased (30% vs 20%), blood sodium decreased (28% vs 20%), lipase increased (25% vs 16%), aspartate aminotransferase (AST) increased (24% vs 12%), blood potassium increased (24% vs 16%), alanine aminotransferase (ALT) increased (24% vs 12%), blood cholesterol increased (22% vs 16%), serum amylase increased (21% vs 12%), hemoglobin decreased (28% vs 18%), and white blood cell decreased (20% vs 10%).

**Fatal adverse reactions** occurred in 1.8% of patients with **advanced renal cell carcinoma (RCC)** receiving BAVENCIO in combination with axitinib. These included sudden cardiac death (1.2%), stroke (0.2%), myocarditis (0.2%), and necrotizing pancreatitis (0.2%).

**The most common adverse reactions** (all grades,  $\geq 20\%$ ) in patients with **advanced RCC** receiving BAVENCIO in combination with axitinib (vs sunitinib) were diarrhea (62% vs 48%), fatigue (53% vs 54%), hypertension (50% vs 36%), musculoskeletal pain (40% vs 33%), nausea (34% vs 39%), mucositis (34% vs 35%), palmar-plantar erythrodysesthesia (33% vs 34%), dysphonia (31% vs 3.2%), decreased appetite (26% vs 29%), hypothyroidism (25% vs 14%), rash (25% vs 16%), hepatotoxicity (24% vs 18%), cough (23% vs 19%), dyspnea (23% vs 16%), abdominal pain (22% vs 19%), and headache (21% vs 16%).

**Selected laboratory abnormalities** (all grades,  $\geq 20\%$ ) worsening from baseline in patients with **advanced RCC** receiving BAVENCIO in combination with axitinib (vs sunitinib) were blood triglycerides increased (71% vs 48%), blood creatinine increased (62% vs 68%), blood cholesterol increased (57% vs 22%), alanine aminotransferase increased (ALT) (50% vs 46%), aspartate aminotransferase increased (AST) (47% vs 57%), blood sodium decreased (38% vs 37%), lipase increased (37% vs 25%), blood potassium increased (35% vs 28%), platelet count decreased (27% vs 80%), blood bilirubin increased (21% vs 23%), and hemoglobin decreased (21% vs 65%).



Please see Important Safety Information on pages 13-15.

Click for full [Prescribing Information and Medication Guide](#), or visit [BAVENCIO.com](https://www.bavencio.com).

**References:** **1.** Centers for Medicare and Medicaid Services CMS, Alpha-Numeric HCPCS File, October 2023. **2.** CMS. CY 2023 Medicare Physician Fee Schedule Final Rule (public display version). November 2, 2022. Page 939. **3.** Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual, Chapter 15. Accessed October 25, 2023. <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf> **4.** CMS. CY 2023 Medicare Hospital Outpatient Prospective Payment Final Rule (public display version). October 25, 2023. **5.** Social Security Act, Section 1847A. Accessed October 25, 2023. [https://www.ssa.gov/OP\\_Home/ssact/title18/1847A.htm](https://www.ssa.gov/OP_Home/ssact/title18/1847A.htm) **6.** CMS 2024 Medicare Hospital Outpatient Prospective Payment Final Rule, CMS-1786-F, November 2023. **7.** CMS, Medicare Part B ASP Files. Accessed October 25, 2023. <https://www.cms.gov/medicare/payment/all-fee-service-providers/medicare-part-b-drug-average-sales-price/asp-pricing-files> **8.** Centers for Medicare and Medicaid Services. Medicare Managed Care Model Chapter 4. Accessed October 25, 2023. <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/mc86c04.pdf>

EMD Serono, Inc. does not guarantee coverage and/or reimbursement for BAVENCIO. Coverage, coding, and reimbursement policies vary significantly by payer, patient, and setting of care. Actual coverage and reimbursement decisions are made by individual payers following the receipt of claims. Patients and healthcare professionals should always verify coverage, coding, and reimbursement guidelines on a payer- and patient-specific basis.